


Can individuals use smoke forecasts for personal decision making? A call to action

Renato Berlinghieri*, David Burt*, Paolo Giani, Arlene Fiore, Tamara Broderick


- A person needs to go out during the day to take care of a sick family member. They are flexible about exactly when and want to limit PM2.5 exposure. Can they use model forecasts to decide when to go out?
- We analyze extreme events and look at cities across the continental US.



Can individuals use smoke forecasts for personal decision making? A call to action

Renato Berlinghieri*, David Burt*, Paolo Giani, Arlene Fiore, Tamara Broderick


- A person needs to go out during the day to take care of a sick family member. They are flexible about exactly when and want to limit PM2.5 exposure. Can they use model forecasts to decide when to go out?
- We analyze extreme events and look at cities across the continental US.



Can individuals use smoke forecasts for personal decision making? A call to action

Renato Berlinghieri*, David Burt*, Paolo Giani, Arlene Fiore, Tamara Broderick


- A person needs to go out during the day to take care of a sick family member. They are flexible about exactly when and want to limit PM2.5 exposure. Can they use model forecasts to decide when to go out?
- We analyze extreme events and look at cities across the continental US.



Can individuals use smoke forecasts for personal decision making? A call to action

Renato Berlinghieri*, David Burt*, Paolo Giani, Arlene Fiore, Tamara Broderick


- A person needs to go out during the day to take care of a sick family member. They are flexible about exactly when and want to limit PM2.5 exposure. Can they use model forecasts to decide when to go out?
- We analyze extreme events and look at cities across the continental US.
- Baseline: predict the level of PM2.5 read at monitors to remain constant over a 24 hour window.



Can individuals use smoke forecasts for personal decision making? A call to action

Renato Berlinghieri*, David Burt*, Paolo Giani, Arlene Fiore, Tamara Broderick

- A person needs to go out during the day to take care of a sick family member. They are flexible about exactly when and want to limit PM2.5 exposure. Can they use model forecasts to decide when to go out?
- We analyze extreme events and look at cities across the continental US.
- Baseline: predict the level of PM2.5 read at monitors to remain constant over a 24 hour window.
- Existing predictive models (NOAA HRRR-Smoke and CAMS) don't provide reliable enough information for this type of decision making.



Can individuals use smoke forecasts for personal decision making? A call to action

Renato Berlinghieri*, David Burt*, Paolo Giani, Arlene Fiore, Tamara Broderick

- A person needs to go out during the day to take care of a sick family member. They are flexible about exactly when and want to limit PM2.5 exposure. Can they use model forecasts to decide when to go out?
- We analyze extreme events and look at cities across the continental US.
- Baseline: predict the level of PM2.5 read at monitors to remain constant over a 24 hour window.
- Existing predictive models (NOAA HRRR-Smoke and CAMS) don't provide reliable enough information for this type of decision making.

